



**UNIVERSITI PUTRA MALAYSIA**

**MICROBIAL AND PHYSICOCHEMICAL CHANGES DURING  
FERMENTATION OF THAI CHICKEN SAUSAGE**

**APINYA VANICHPUN**

**FSMB 2003 23**

**MICROBIAL AND PHYSICOCHEMICAL CHANGES DURING  
FERMENTATION OF THAI CHICKEN SAUSAGE**

**APINYA VANICHPUN**

**MASTER OF SCIENCE  
UNIVERSITI PUTRA MALAYSIA**

**2003**



**MICROBIAL AND PHYSICOCHEMICAL CHANGES DURING  
FERMENTATION OF THAI CHICKEN SAUSAGE**

**By**

**APINYA VANICHPUN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
In Fulfilment of the Requirements for the Degree of Master of Science**

**August 2003**



Dedicated to

My family for their loves and supports and special dedication to Lee Yock

Ann who warmed my heart while I was away from home

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Science

**MICROBIAL AND PHYSICOCHEMICAL CHANGES DURING  
FERMENTATION OF THAI FERMENTED CHICKEN SAUSAGE**

**By**

**APINYA VANICHPUN**

**August 2003**

**Chairman : Dr. Zaiton Hassan, Ph.D.**

**Faculty : Food Science and Biotechnology**

A newly formulated fermented Thai chicken sausage was developed by mixing chicken meat with glutinous rice, sugar, garlic, salt, and black pepper. Glutinous rice rendered good binding of the mixed ingredients when compared to other carbohydrate sources (rice, glass noodle, rice + glass noodle, and glutinous rice + glass noodle). Moreover, glutinous rice also significantly ( $p < 0.05$ ) reduced the pH value from 6.06 to 4.42 when compared to the formulation with added rice (from 5.80 to 4.50), rice + glass noodle (from 5.77 to 4.43), and glutinous rice + glass noodle (from 5.82 to 4.34) within 24h of fermentation. The addition of water (5% w/w) with and without sodium tripolyphosphate (0.2% w/w) significantly ( $p < 0.05$ ) improved moisture retention in the sausages. Sodium tripolyphosphate (STPP) and water significantly ( $p < 0.05$ ) affect the pH reduction and the growth of lactic acid bacteria. The formation of mainly lactic acid from 4.96 to 23 mmol of acid/ 100 g of samples and a small amount of acetic acid from 1.57 to 3.27 mmol of acid/ 100 g of samples as a result of

carbohydrate fermentation by LAB during 0h-48h fermentation contributed to the acidic aroma and typical tangy taste typically found in Thai fermented sausages. The fermentation process was successfully carried out within 48h as shown by the rapid increase in the growth of LAB from 7.80  $\log_{10}\text{cfu/g}$  to 12.46  $\log_{10}\text{cfu/g}$ , rapid increase in titratable acidity from 0.90 to 2.25 g of lactic acid/ 100 g of sausages, and sharp decrease in pH values from 5.64 to 4.27. Prolonged fermentation (after 48h) caused an over acidity in the sausages with a final pH between 4.20-4.30. This newly formulated fermented chicken sausage proved to be highly selective for the growth of LAB. Four species of LAB namely, *Pediococcus pentosaceus*, *Pd. acidilactici*, *Lactobacillus pentosus*, and *Lb. brevis* were isolated. *Pd. pentosaceus* and *Pd. acidilactici* were found throughout the fermentation process, whereas *Lb. pentosus* and *Lb. brevis* were found after 24h of fermentation, indicating that the latter two species are tolerant to the acidic conditions. *Pd. pentosaceus* was found throughout the fermentation process in higher number than other species.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
Sebagai memenuhi keperluan untuk ijazah Master Sains

**PERUBAHAN MIKROBIOLOGI DAN FISIKO-KIMIA SEMASA  
FERMENTASI SOSEJ AYAM THAI**

Oleh

**APINYA VANICHPUN**

**Ogos 2003**

**Pengerusi: Dr. Zaiton Hassan, Ph.D.**

**Fakulti: Sains Makanan dan Bioteknologi**

Formulasi baru sosej ayam Thai telah dihasilkan mengandungi daging ayam cincang yang dicampur dengan pulut masak, bawang putih, gula, garam dan lada hitam. Pulut memberikan kapasiti keupayaan pengabungan bahan-bahan campuran yang baik berbanding karbohidrat lain (nasi, tanghun, nasi + tanghun, dan pulut + tanghun). Tambahan pula, pulut memberikan signifikasi penurunan nilai pH dari 6.06 kepada 4.42 ( $p < 0.05$ ) berbanding formulasi yang ditambahkan nasi (dari 5.80 kepada 4.50), nasi + tanghun (dari 5.77 kepada 4.43), atau nasi pulut + tanghun (dari 5.82 kepada 4.34) dalam fermentasi 24 jam. Penambahan air (5% w/w) dengan atau tanpa natrium trifosfat (0.2% w/w) memberikan signifikasi ( $p < 0.05$ ) pengekalan kelembapan di dalam sosej. STPP dan air memberi kesan signifikan pH penurunan ( $p < 0.05$ ). Pembentukan asid utama melibatkan asid laktik dari 4.96 kepada 14.23 mmol asid/100g sampel) dan jumlah asid asetik yang kecil dari 1.57 kepada 3.27 mmol asid /100g sampel sebagai hasil fermentasi karbohidrat oleh LAB semasa fermentasi

0-48 jam turut menyumbangkan aroma asidik dan rasa masam yang biasa ditemui dalam sosej fermentasi Thai. Proses fermentasi telah dijalankan dengan dalam masa 48 jam menunjukkan pertumbuhan pantas di dalam LAB dengan peningkatan bilangan LAB dari  $\log_{10} 7.80$  cfu/g kepada  $\log_{10} 12.46$  cfu/g, titrasi asiditi dari 0.90 kepada 2.25 asid laktik /100 g sosej, dan penurunan nilai pH dari 5.64 kepada 4.27. Fermentasi berpanjangan (selepas 48 jam) menyebabkan asiditi berlebihan di dalam sosej dengan pH akhir 4.20-4.30. Formulasi sosej ayam yang baru ini telah memberikan pemilihan yang tinggi untuk pertumbuhan LAB. Empat jenis LAB yang telah dipencilkan adalah *Pediococcus pentosaceus*, *Pd. acidilactici*, *Lactobacillus pentosus*, dan *Lb. brevis*. *Pd. pentosaceus* dan *Pd. acidilactici* telah ditemui pada setiap peringkat fermentasi, manakala, *Lb. pentosus* dan *Lb. brevis* ditemui selepas fermentasi 24 jam. Ini menunjukkan dua spesis terakhir adalah toleran kepada keadaan asidik. *Pd. pentosaceus* ditemui di sepanjang proses fermentasi dan mempunyai bilangan jumlah yang jauh lebih tinggi berbanding spesis lain.



## ACKNOWLEDGEMENTS

I would like to express my most sincere gratitude to the Southeast Asian Ministers of Education Organization REGIONAL CENTER FOR GRADUATE STUDY AND RESEARCH IN AGRICULTURE (SEARCA) for supporting me to pursue my Masters degree study at Universiti Putra Malaysia. My sincere gratitude is also extended to my home institution “Rajamangala Institute of Technology” Ministry of Education, Thailand for granting me study leave.

I wish to express my sincere gratitude and appreciation to my supervisor, Dr. Zaiton Hassan for her guidance, advice, and comment throughout the course of my study. Appreciation is also due to the members of my supervisory committee, Associate Professor Dr. Jamilah Bakar, and Dr. Nazimah Sheikh Abdul Hamid for their advices and suggestions.

Appreciations are also extended to all members of the Faculty of Food Science and Biotechnology, UPM for providing the research facilities and technical assistance during my graduate study. Acknowledgement is also due to all my friends, especially Lee Yock Ann for her kind help, suggestion, and technical assistance, Narumon Sumalee, Wanna Ammawath, Pitoon Nopnakom, Benchamaporn Pimpa, Bodee Khamseekieow, Youssr Abdul Hadi Noaman, Noor Zaleha Bt. Awang Salleh, Ooi Wai Ling, Lim Yin Sze, Samuel Lihan, Ahmad Zainuri Dzomir, Siti



Zulaiha Bt. Hanapi, Sirinath Srionnuan, Assistant Professor Suwit Suwanno, Assistant Professor Dr. Sarakam Kaewtasi, Jamnanporn Choykeow, Cheunsikah Chemong, and Saowanee Chaipetch, who had given me the moral encouragement and support to complete my graduate study.

I certify that an Examination Committee met on 12<sup>nd</sup> August 2003 to conduct the final examination of Apinya Vanichpun on her Master of Science thesis entitled "Microbial and Physicochemical Changes during Fermentation of Thai Chicken Sausage" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Putra Malaysia (Higher Degree) Regulations 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**GULAM RUSUL RAHMAT ALI, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Chairman)

**ZAITON HASSAN, Ph.D.**

Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**JAMILAH BAKAR, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**NAZIMAH SHEIKH ABDUL HAMID, Ph.D.**

Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)



---

**GULAM RUSUL RAHMAT ALI, Ph.D.**  
Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 04 DEC 2003

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science

**ZAITON HASSAN, Ph.D.**

Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Chairman)

**JAMILAH BAKAR, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**NAZIMAH SHEIKH ABDUL HAMID, Ph.D.**

Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)



**AINI IDERIS, Ph.D.**

Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 8 JAN 2024

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

Apinya Vanichpun

APINYA VANICHPUN

Date: 2 DEC 2003

## TABLE OF CONTENTS

DEDICATION.....	ii
ABSTRACT.....	iii
ABSTRAK.....	v
ACKNOWLEDGEMENTS.....	vii
APPROVAL.....	ix
DECLARATION.....	xi
LIST OF TABLES.....	xv
LIST OF FIGURES.....	xix
LIST OF PLATES.....	xx
LIST OF ABBREVIATIONS.....	xxi
 <b>CHAPTER</b>	
<b>I     GENERAL INTRODUCTION.....</b>	<b>1</b>
 <b>II    LITERATURE REVIEW.....</b>	<b>4</b>
Classification of Sausages.....	5
Ingredients for Sausage Production.....	5
Meat.....	5
Fat.....	6
Carbohydrates.....	6
Salt.....	10
Spices.....	11
Phosphates.....	12
Nitrite/Nitrate.....	14
Sausage Production.....	17
Comminuting/Grinding/Chopping/Mincing.....	18
Stuffing.....	18
Casing.....	20
Fermentation.....	21
Smoking.....	22
Fermented Sausages.....	24
Definition .....	24
Fermented Sausages Classification.....	25
Western Fermented Sausages.....	25
Eastern Fermented Sausages.....	27
Microorganisms Involved in Fermented Sausages	
Production.....	32
Lactic Acid Bacteria.....	34
Homofermentative and Heterofermentative Lactic acid	
Bacteria .....	35
Lactic Acid Bacteria and Acidification Process.....	37
<i>Micrococcaceae</i> and <i>Staphylococcaceae</i> .....	38
Moulds and Yeasts.....	42

	Flavour Development.....	44
	Proteolysis.....	44
	Lipolysis.....	46
	Formation of Biogenic Amines.....	48
	Starter Cultures Used in Fermented Sausages production.....	50
<b>III</b>	<b>OPTIMIZATION OF THE FORMULATION FOR CHICKEN SAUSAGE.....</b>	<b>55</b>
	Introduction.....	55
	Materials and Methods.....	56
	Materials.....	56
	Experimental Design.....	56
	Chemical Analysis.....	64
	Microbiological Analysis.....	66
	Statistical Analysis.....	67
	Results.....	67
	Discussion.....	78
	Conclusion.....	82
<b>IV</b>	<b>THE EFFECTS OF SODIUM TRIPOLYPHOSPHATE AND WATER ADDITION ON CHEMICAL AND MICROBIOLOGICAL CHANGES DURING FERMENTATION.....</b>	<b>83</b>
	Introduction.....	83
	Materials and Methods.....	84
	Materials.....	84
	Experimental Design.....	84
	Chemical Analysis.....	85
	Microbiological Analysis.....	87
	Statistical Analysis.....	87
	Results.....	87
	Discussion.....	95
	Conclusion.....	107
<b>V</b>	<b>ISOLATION AND IDENTIFICATION OF LACTIC ACID BACTERIA FROM THE NEWLY FORMULATED FERMENTED CHICKEN SAUSAGE.....</b>	<b>109</b>
	Introduction.....	109
	Materials and Methods.....	110
	Media and Cultures Conditions.....	110
	Physiological and Biochemical Tests.....	111
	Tests Reproducibility.....	113
	Results.....	115
	Discussion.....	136
	Conclusion.....	150
<b>VI</b>	<b>GENERAL DISCUSSION.....</b>	<b>151</b>

<b>VII CONCLUSION AND RECOMMENDATIONS.....</b>	<b>161</b>
BIBLIOGRAPHY.....	165
APPENDICES.....	181
BIODATA OF THE AUTHOR.....	212



## LIST OF TABLES

Table	Page
2.1 Classification of sausages.....	8
2.2 Characteristics of dry and semi-dry sausages.....	26
2.3 Processing features of Western fermented sausages.....	28
2.4 Lactic acid bacteria strains and other organisms isolated from fermented sausage.....	33
2.5 Factors affecting the acidification process during sausage fermentation.....	40
2.6 Biogenic amines production by the decarboxylases microorganisms.....	51
2.7 A list of microorganisms used as a starter culture in fermented sausages production.....	53
3.1 Formulation of Thai fermented chicken sausage containing different complex carbohydrate sources.....	61
3.2 Formulation of Thai fermented chicken sausage containing different ratio of glutinous rice to sugar.....	62
3.3 Formulation of Thai fermented chicken sausage containing different complex carbohydrate sources to STPP.....	63
3.4 Formulation of Thai fermented chicken sausage containing different amount of sugar in the presence or absence of STPP and water.....	65
3.5 Changes in pH of sausages containing different complex carbohydrate sources during fermentation.....	68
3.6 Changes in pH of sausages made of different concentration of glutinous rice and sugar.....	69
3.7 Changes in titratable acidity (g of lactic acid / 100 g of samples) of sausages made of different concentration of glutinous rice and sugar.....	71

3.8	Changes in moisture content (%) of sausages made of different concentration of glutinous rice and sugar.....	72
3.9	Sensory properties of sausages of different concentration of glutinous rice and sugar after 48h fermentation.....	74
3.10	Changes in pH and moisture content (% wet weight) of sausages made of different carbohydrate sources in the presence or absence of sodium tripolyphosphate.....	75
3.11	Changes in titratable acidity (g of lactic acid/ 100 g of samples) pH, and moisture content (%) of sausages made of different concentration of sugar in the presence or absence of STPP and water.....	77
3.12	Changes in total viable counts (TVC) and lactic acid bacteria counts (LAB) ( $\log_{10}$ cfu/g) of sausages made of different concentration of sugar in the presence or absence of STPP and water.....	78
4.1	Formulations of ingredients used in this study.....	84
4.2	Changes in pH and titratable acidity (g of lactic acid/ 100 g of samples) of sausages made of different ratio between chicken meat to glutinous rice in the presence or absence of STPP and/or water.....	89
4.3	Changes in moisture content (% wet weight) of sausages made of different ratio between chicken meat to glutinous rice in the presence or absence of STPP and/or water.....	90
4.4	Changes in organic acid profiles during sausage fermentation.....	93
4.5	Changes in total viable counts (TVC) and total lactic acid bacteria counts (LAB) ( $\log_{10}$ cfu/g) of sausages made of different ratio between chicken meat to glutinous rice in the presence or absence of STPP and/or water.....	94
4.6	Scores obtained from sensory evaluation of fermented chicken sausages made of different ratio between chicken meat to glutinous rice in the presence or absence of STPP and/or water.....	96

5.1	Differential characteristics of lactic acid bacteria.....	114
5.2	The numbers of cocci in pairs and/or chains forming LAB after differentiated into subgroup.....	119
5.3	Grouping of cocci in pair and/or chain bacteria isolated from fermented chicken sausages.....	120
5.4	The numbers of rods in pairs and/or chains forming LAB after differentiated into subgroup.....	122
5.5	Grouping of cocci in pair and tetrad bacteria isolated from fermented chicken sausage.....	123
5.6	The numbers of cocci in pairs and tetrads forming LAB after differentiated into subgroup.....	125
5.7	Grouping of <i>Pediococcus</i> sp. strains isolated from fermented chicken sausages.....	126
5.8	Characteristics differentiating of <i>Pediococcus</i> sp.....	130
5.9	The numbers of cocci in pairs and tetrads forming LAB after differentiated into subgroup using the Bergey's Manual Determinative of Bacteriology.....	132
5.10	Grouping of <i>Pediococcus</i> sp. strains isolated from fermented chicken sausages using the Bergey's Manual Determinative of Bacteriology.....	134
5.11	Metabolic profiles of the representative isolate belonging to species <i>Pediococcus pentosaceus</i> 1 using the API 50 CH System.....	138
5.12	Metabolic profiles of the representative isolate belonging to species <i>Pediococcus pentosaceus</i> 2 using the API 50 CH System.....	140
5.13	Metabolic profiles of the representative isolate belonging to species <i>Pediococcus acidilactici</i> using the API 50 CH System.....	143
5.14	Metabolic profiles of the representative isolate belonging to species <i>Lactobacillus pentosus</i> using the API 50 CH System..	146

5.15	Metabolic profiles of the representative isolate belonging to species <i>Lactobacillus brevis</i> 1 using the API 50 CH System.....	148
6.1	The distribution of LAB species found in Thai fermented chicken sausage during fermentation.....	152
6.2	The distribution of LAB species isolated from traditional Thai fermented meat.....	152
6.3	Optimum temperature for growth of LAB species.....	155
6.4	Optimum pH for growth of LAB species.....	155

## LIST OF FIGURES

Figure		Page
2.1	Development of colour in dry sausage.....	16
2.2	Processing of emulsion-type sausage.....	19
2.3	Processing of non emulsion-type sausage.....	19
2.4	Glycolytic pathway in lactic acid bacteria.....	36

## LIST OF PLATES

Plate		Page
3.1	Placing the ingredients into a silent cutter.....	59
3.2	Mixing the mixed ingredients for 1 min using a silent cutter.....	59
3.3	Transferring the mixed ingredients into a stuffer.....	60
3.4	Filling the mixed ingredients into a cellulose casing.....	60
3.5	Fermentation of the stuffed sausages in the incubator at 30°C for 72h.....	61
4.1	Scanning electron micrograph (20x magnification) of fermented chicken sausage after 24h fermentation.....	101
4.2	Scanning electron micrograph (20x magnification) of fermented chicken sausage after 72h fermentation.....	101
5.1	The colonies of lactic acid bacteria on MRS agar plate after incubation at 30°C for 48h.....	112
5.2	Checking the purity of LAB isolate using streak plate technique on MRS agar plate after incubation for 48h.....	112
5.3	Cocci in pair and/or chain bacteria isolated from fermented chicken sausage.....	117
5.4	Rods in pair bacteria isolated from fermented chicken sausage.....	117
5.5	Cocci in pair and tetrad bacteria isolated from fermented chicken sausage.....	118
5.6	The API 50 CH strip of isolate No. 52420 (subgroup 2.6) after incubation at 30°C for 48 h ( <i>Lactobacillus pentosus</i> with 99.8% similarity).....	129
5.7	The API 50 CH strip of the isolate No. 7214 (subgroup 2.13) after incubation at 30°C for 48 h ( <i>Lactobacillus brevis</i> 1 with 84.7% similarity).....	129

## LIST OF ABBREVIATIONS

ADP	= Adenosine-5'-diphosphate
ANOVA	= Analysis of variance
AOAC	= Association of Official Analytical Chemists
ATP	= Adenosine-5'-triphosphate
B.C	= Before Christ
°C	= Degree celcius
CO <sub>2</sub>	= Carbondioxide
DNA	= Deoxyribonucleic acid
EMP	= Embden Meyerhof Parnas
FDP	= Fructose diphosphate aldolase
G+C	= Guanine + Cytosine
g	= Gram
HPLC	= High Performance Liquid Chromatography
H <sub>2</sub> O	= Water
H <sub>2</sub> O <sub>2</sub>	= Hydrogen Peroxide
h	= Hour
kg	= Kilogram
LAB	= Lactic Acid Bacteria
LDH	= Lactate dehydrogenes
Min.	= Minute
mmol	= Millimolar
MRS	= de Man Ragosa Shape

## CHAPTER I

### GENERAL INTRODUCTION

Sausages are popular meat products enjoyed by millions of consumer worldwide (Mendoza *et al.*, 2001). In Thailand, the total consumption of sausage in 1988 was 5,000 tons and this include 1,000 ton of fermented sausage, however the total production of fermented sausage by Sor Khonkean, a Thai company that produces meat products reached up to 9,600 tons/year in 1993 (Sartsittisuk, 1995). In the United States of America, over 2.18 billion kilograms of sausages are produced annually (Matulis *et al.*, 1995). In Germany, a large variety of fermented sausages have been developed with over 350 varieties being produced (Vernam and Sutherland, 1995). The annual per capita consumption of fermented sausages in Germany is in the order of 5 kg (Lücke, 1985). About 50 varieties of fermented sausages have been catalogued in Spain (Fernández *et al.*, 2000) and the production of *Chorizo*, the main fermented sausage produced in Spain was 61,700 tons annually (Pagan-Moreno *et al.*, 1998).

Lactic acid bacteria (LAB) are important in the ripening process of fermented sausages (Hammes *et al.*, 1994), and they are usually the predominant microbial population in this type of sausage (Mäkelä, 1992; Bruna *et al.*, 2001a; Coppola *et al.*, 2000). This is because the production of



organic acids, mainly lactic acid and low concentrations of acetic acid by LAB result in a decrease in pH, which is responsible for the preservative effect. The combined effect between lactic and acetic acid imparts an “acid flavour” which is predominant in semi-dry sausages (Lücke, 2000; Vignolo *et al.*, 1989). In addition, the accumulation of lactic and acetic acid can inhibit meat-borne pathogenic bacteria, coagulate soluble meat proteins, and facilitating drying of the products (Lücke, 2000).

Throughout the industrialized world, a growing concern for health has increased the interest in protein sources that are low in fat and cholesterol (Pedersen, 1995). The use of chicken meat as a raw material for different chicken meat based products is becoming popular (Silva and Glória, 2002). Currently, health-conscious consumers try to reduce dietary fat intake by consuming low fat, reduced fat or fat free foods. This has in turn increased consumer demand for low fat content meat products (Mugumura *et al.*, 2003) and resulted in the development of new formulations or modification of traditional food products to contain less fat (Mendoza *et al.*, 2001).

The traditional Thai fermented sausage is mainly made up of ground lean pork mixed with lard. This resulted in a high fat content of the final product. Thungtakul (1988) reported that the content of fat in Thai fermented sausage was 38.98% (dry basis) after 144h of fermentation.